

# We'll Give You a Hand: Wrist & Hand Pathologies You May Be Missing

Jerry Weniger, PhD, ATC, PA-C

Director, Physician Assistant Program  
Associate Professor, Health Professions, Dept.  
James Madison University

# DISCLOSURES

I have no personal or financial interests to declare.

I receive no financial support from industry sources.

Images were taken from fair use and creative commons libraries.



# Elbow special tests (we'll come back to these later)



Hand/wrist special tests (we'll come back to these later)



# PRE-TEST QUESTION #1

- ◉ Why is it important to diagnose Kienböck's disease as early as possible?
  - A. The disease course can be reversed with early pharmacologic intervention
  - B. To prevent the spread of necrosis to adjacent bones
  - C. Because ultrasound can be both diagnostic and therapeutic for the condition
  - D. To intervene before bony collapse of the lunate occurs

# PRE-TEST QUESTION #2

- ⦿ Definitive treatment for a Jersey finger injury...
  - A. is always conservative: 6-8 weeks of splinting typically does well.
  - B. may be conservative or surgical, it depends on the location of the injury.
  - C. is always surgical (primary tendon repair or fracture fragment repair). Long-term splinting is rarely an option.
  - D. is a corticosteroid injection at the site of injury.

# PRE-TEST QUESTION #3

- ◉ When evaluating a patient with a suspected skier's thumb injury...
  - A. it is best to obtain radiographs prior to assessing the UCL.
  - B. radiographs are not necessary - it is a clinical diagnosis.
  - C. it is best to obtain radiographs after assessing the UCL.
  - D. MRI is the gold standard imaging that is needed.


# INTRODUCTION & BACKGROUND

- ⦿ hand & wrist susceptible to injury & overuse
- ⦿ hand function abnormal = disability
- ⦿ ~11.3% of all ED visits in the US involved injuries to the hand, wrist, or fingers



# KIENBÖCK'S DISEASE

- avascular necrosis of lunate
  - leads to progressive collapse
  
- etiology unknown
  - disruption of blood supply
  - undiagnosed fracture?
  - repetitive trauma?




# KIENBÖCK'S DISEASE

- more common in males

- dorsal wrist pain

- vague complaints

- wrist swelling
- wrist stiffness



# KIENBÖCK'S DISEASE

- over time...

- crepitus
- ↓ ROM
- weakness with grip




- progression varies

- typically over several years

# KIENBÖCK'S DISEASE

## ○ Radiographs

- shows increased density of lunate
- not very sensitive



# KIENBÖCK'S DISEASE


- More sensitive imaging, helpful for early disease
  - MRI
    - decreased signal on T1 image



# KIENBÖCK'S DISEASE

## Radiograph considerations:

- “ulna positive variance” vs “ulna negative variance”



# KIENBÖCK'S DISEASE




Image from UpToDate © 2018

# KIENBÖCK'S DISEASE

## Treatment (conservative):

- immobilization





# KIENBÖCK'S DISEASE

## Treatment (surgical):



- 1<sup>st</sup> line options:
  - radial shortening osteotomy
  - vascularized bone graft
- “salvage procedures”:
  - proximal row carpectomy
  - wrist arthrodesis



# COLLATERAL LIGAMENT INJURIES

## UCL Tear (MCL)

- valgus force
- overhead throwing athletes!
  - especially pitchers
  - late cocking, early acceleration
  - *loss of velocity/accuracy*



# COLLATERAL LIGAMENT INJURIES

## UCL Tear Prevention

### Little League Pitch Counts

Age	Daily Max Pitches	Required Rest Days (Pitches)				
		0 Days	1 Day	2 Days	3 Days	4 Days
7-8	50	1-20	21-35	36-50	N/A	N/A
9-10	75	1-20	21-35	36-50	51-65	66+
11-12	85	1-20	21-35	36-50	51-65	66+
13-14	95	1-20	21-35	36-50	51-65	66+
15-16	95	1-30	31-45	46-60	61-75	76+
17-18	105	1-30	31-45	46-60	61-75	76+

Source: <https://www.littleleague.org/playing-rules/pitch-count/>

# COLLATERAL LIGAMENT INJURIES

## UCL Tear

- special tests



Valgus Stress Test




Milking Maneuver

# COLLATERAL LIGAMENT INJURIES

## UCL Tear

- MRI Arthrogram






# COLLATERAL LIGAMENT INJURIES





## UCL Tear

- no conservative treatment
- surgery: UCL reconstruction
  - “Tommy John”



# DEQUERVAIN'S TENOSYNOVITIS



- 1<sup>st</sup> dorsal extensor compartment
  - abductor pollicis longus
  - extensor pollicis brevis
- pain with lifting (“new mommy syndrome”)



# DEQUERVAIN'S TENOSYNOVITIS

- TTP along tendons near radial styloid
- “snowball crepitus”
- pain with...
  - resistive thumb extension
  - resistive ulnar deviation of wrist

- Special Test






# DEQUERVAIN'S TENOSYNOVITIS

- typically no imaging necessary



- Treatment

- NSAIDS, RICE
- thumb spica splint
- PT/OT referral
- corticosteroid injection



Inject at an angle of 15 to 20 degrees in line with abductor pollicis longus tendon, approximately 1 cm distal to the point of maximal tenderness and directed proximally.

# GANGLION CYST

- most common soft tissue mass of hand/wrist
  - some are painful, not always
- more common in women, more common dorsally
  - 70% are dorsal, near scaphoid or lunate
- weakness in tendon sheath capsule
  - previous trauma?



# GANGLION CYST

## ○ clinical diagnosis


- +/- Xrays
- trans-illumination
- ultrasound



# GANGLION CYST

## ○ Treatment

- splint
- aspiration
- *surgical excision*




“Bible Bump”

# MALLET FINGER



# MALLET FINGER

- sudden flexion force
  - typically from object




- causes flexion deformity/extensor lag at the DIP




# MALLET FINGER

- injury to extensor mechanism @ dorsal DIP joint
  - may be tendon rupture
  - may be avulsion fracture



# MALLET FINGER

- ◉ ecchymosis, swelling over DIP
- ◉ TTP at distal finger (DIP), especially dorsal
- ◉ flexion deformity/extensor lag
- ◉ pain with motion





# MALLET FINGER

- finger X-rays (not hand)
  - AP, lateral, oblique



# MALLET FINGER

## ○ Soft Tissue Mallet

- 6-8 weeks of extension splinting
- may initiate within 3 months of injury



# MALLET FINGER

- Soft Tissue Mallet

- 6-8 weeks of extension splinting




- do not immobilize PIP

# MALLET FINGER

## ○ Soft Tissue Mallet



- if conservative treatment fails...




# MALLET FINGER

## ○ Bony Mallet (fracture)

- treat with 6-8 weeks of extension splinting unless...
  - fracture fragment > 50% articular surface
  - dislocation with fracture



# JERSEY FINGER



# JERSEY FINGER

- ⦿ sudden hyperextension of DIP *during active flexion*
  - caught in shirt/jersey
  - football
  
- ⦿ ring finger most common



# JERSEY FINGER

- injury to FDP tendon @ volar distal phalanx
  - may be tendon rupture
  - may be avulsion fracture





Image from UpToDate © 2018




# JERSEY FINGER

- ecchymosis, swelling over volar finger
- TTP at distal finger, especially volar
- slight flexion deformity
- *may* palpate lump in palm





# JERSEY FINGER

- cannot flex the **DIP** (yet can still flex PIP)
  - must evaluate DIP flexion in isolation!



# JERSEY FINGER

- cannot flex the **DIP** (yet can still flex PIP)
  - must evaluate DIP flexion in isolation!



# JERSEY FINGER

- ◉ finger X-rays (not hand)
  - AP, lateral, oblique  
(bony jersey finger not as common as bony mallet finger)
- ◉ Initially, splint in flexion
  - “extension block splint”



# JERSEY FINGER

- typically no conservative treatment
  - splinting (long term) *rarely* an option
- surgery
  - primary tendon repair
  - fracture fragment repair



# GAMEKEEPER'S THUMB - SKIER'S THUMB



# GAMEKEEPER'S THUMB - SKIER'S THUMB

- first recognized in Scottish “gamekeepers”
  - repetitive “neck wringing’ of game between thumb & index finger - “gamekeeper’s thumb”



# GAMEKEEPER'S THUMB - SKIER'S THUMB

- ◉ Skier's thumb
  - from acute injury, usually a fall





# GAMEKEEPER'S THUMB - SKIER'S THUMB

- ◉ Gamekeeper's Thumb: chronic, overuse
- ◉ Skier's Thumb: acute injury





Image from UpToDate © 2018

# GAMEKEEPER'S THUMB - SKIER'S THUMB

- injury to thumb ulnar collateral ligament (UCL)
  - base of the proximal phalanx at the 1<sup>st</sup> MCP
  - either partial or complete tear
  - with or without fracture
  - acute or chronic




# GAMEKEEPER'S THUMB - SKIER'S THUMB

- valgus & hyperextension force to thumb

- common injury

- skiers
- football lineman
- any FOOSH




# GAMEKEEPER'S THUMB - SKIER'S THUMB

- ⦿ “jammed thumb”
- ⦿ pain, swelling at 1<sup>st</sup> MCP
- ⦿ ecchymosis, thenar eminence
- ⦿ painful ROM



# GAMEKEEPER'S THUMB - SKIER'S THUMB

- ⦿ do not stress MCP joint prior to X-rays!
  - must r/o fracture first
  - do not want to displace bony fragment




# GAMEKEEPER'S THUMB - SKIER'S THUMB

- If fracture is present; orthopedic referral...
  - do not stress the ligament during physical exam




# GAMEKEEPER'S THUMB - SKIER'S THUMB

- If fracture ruled out...
- Physical exam: valgus stress
  - increased laxity? definitive endpoint?
  - compare to other side



# GAMEKEEPER'S THUMB - SKIER'S THUMB

- If physical exam is equivocal...and standard radiographs have already demonstrated no fracture:
  - stress radiographs
  - MRI may be necessary
  - ultrasound?





# GAMEKEEPER'S THUMB - SKIER'S THUMB

## ○ Initial Management

- thumb spica splint
- refer to Orthopedics




Image from UpToDate © 2018

# GAMEKEEPER'S THUMB - SKIER'S THUMB

- Definitive Management

- partial tear or non-displaced fracture: cast/splint



Image from UpToDate © 2018




Image from UpToDate © 2018

# GAMEKEEPER'S THUMB - SKIER'S THUMB


## ○ Definitive Management

- partial tear or non-displaced fracture: cast/splint
- complete tear or displaced fracture: surgery




# CARPAL TUNNEL SYNDROME

- elevated pressure within carpal tunnel puts pressure on median nerve
  - short term = numbness and tingling in the fingers
  - long term = nerve damage and muscle weakness



# CARPAL TUNNEL SYNDROME

- elevated pressure within carpal tunnel puts pressure on median nerve
  - short term = numbness and tingling in the fingers
  - long term = nerve damage and muscle weakness



# CARPAL TUNNEL SYNDROME

## ○ more likely in:


- overweight/obese
- females > males
- occupation w/ repetitive hand use
- pregnancy



# CARPAL TUNNEL SYNDROME

## ○ symptoms

- pain & paresthesia's in lateral 3 ½ digits
- worse at night/sleep
- worse with driving
- worse with repetitive activities
- clumsiness with hand



# CARPAL TUNNEL SYNDROME

## Physical Exam:

- muscle loss (late finding)
- muscle weakness (late finding)
- Tinel Sign
- Phalen Test\*
- ***Carpal Compression Test\****




Image from UpToDate © 2018



Image from UpToDate © 2018




# CARPAL TUNNEL SYNDROME

- clinical diagnosis: no imaging studies needed
  - EMG and/or NCV – only if diagnosis is not clear



# CARPAL TUNNEL SYNDROME

- clinical diagnosis: no imaging studies needed
  - bedside ultrasound?



Images from UpToDate © 2018

# CARPAL TUNNEL SYNDROME

- ◉ *Clinical* grading of severity:

# CARPAL TUNNEL SYNDROME

- ◉ *Clinical* grading of severity:

- **Mild:**

- **subjective** symptoms: numbness, tingling, discomfort
- no **sleep** disruption
- not affecting **ADLs**

# CARPAL TUNNEL SYNDROME

## ○ *Clinical* grading of severity:

### ■ **Mild:**

- **subjective** symptoms: numbness, tingling, discomfort
- no **sleep** disruption
- not affecting **ADLs**

### ■ **Moderate:**

- **objective sensory loss** in median nerve distribution
- occasional **sleep** disruption
- symptoms that interfere w/ function but do not limit **ADLs**

# CARPAL TUNNEL SYNDROME

## ◉ *Clinical* grading of severity:

### ■ **Mild:**

- **subjective** symptoms: numbness, tingling, discomfort
- no **sleep** disruption
- not affecting **ADLs**

### ■ **Moderate:**

- **objective sensory loss** in median nerve distribution
- occasional **sleep** disruption
- symptoms that interfere w/ function but do not limit **ADLs**

### ■ **Severe:**

- **weakness** in median nerve distribution
- routinely disrupt **sleep**
- symptoms prevent/limit **ADLs**

# CARPAL TUNNEL SYNDROME

## Conservative Management



- rest
- activity restriction
- NSAIDS
- bracing (daytime vs. nocturnal)
- therapy (PT/OT) to improve function, strength
  - nerve gliding
  - therapeutic ultrasound
  - carpal bone mobilizations



# CARPAL TUNNEL SYNDROME

## Conservative Management

- CS injection
- 80% get relief initially
- Duration of relief?
- No more than 2x per year?





# CARPAL TUNNEL SYNDROME

## Surgical Management

- open or endoscopic
- earlier return to work/activity w/ endoscopic
- no difference in long term outcomes & complications




# TRIGGER FINGER



# TRIGGER FINGER

- “stenosing flexor tenosynovitis”
  - thickened “pulley”
  - more likely in 2, 3, 4<sup>th</sup> fingers
  - worse morning & night
  - palpable nodule



Images from UpToDate © 2018

# TRIGGER FINGER

- limited use for conservative treatment
  - NSAIDS
  - bracing



Image from UpToDate © 2018

# TRIGGER FINGER

## ○ Corticosteroid Injection

- persistent relief beyond 1 year, in 50% of patients
- other 50% may need 2<sup>nd</sup> injection




Image from UpToDate © 2018

# TRIGGER FINGER

## ○ Surgical Release

- 94% success rate overall
- 87% in diabetics



# SUMMARY

## Special Tests

Valgus Stress (elbow)	UCL Tear
Milking Maneuver	
Finkelstein's Test	DeQuervain's Tenosynovitis
Valgus Stress (thumb)	Gamekeeper/Skier Thumb
Tinel Sign	Carpal Tunnel Syndrome
Phalen Sign	
Carpal Compression Test	

# LESSONS FOR PRACTICE

- ◉ **Keinböck's:** MRI needed to see early density changes
- ◉ **UCL Tear:** present when they notice a loss of velocity/accuracy
- ◉ **DeQuervain's:** snowball crepitus, Finkelstein test
- ◉ **Ganglion cyst:** trans-illumination, ultrasound
- ◉ **Mallet finger:** extension splint, +/- surgery
- ◉ **Jersey finger:** check DIP flexion, all will need surgery
- ◉ **Gamekeeper's/Skier's:** no ligamentous testing until after X-rays
- ◉ **Carpal tunnel:** clinical grading of severity



# POST-TEST QUESTION #1

- ◉ Why is it important to diagnose Kienböck's disease as early as possible?
  - A. The disease course can be reversed with early pharmacologic intervention
  - B. To prevent the spread of necrosis to adjacent bones
  - C. Because ultrasound can be both diagnostic and therapeutic for the condition
  - D. To intervene before bony collapse of the lunate occurs

# POST-TEST QUESTION #1

- ◉ Why is it important to diagnose Kienböck's disease as early as possible?
  - A. The disease course can be reversed with early pharmacologic intervention
  - B. To prevent the spread of necrosis to adjacent bones
  - C. Because ultrasound can be both diagnostic and therapeutic for the condition
  - D. *To intervene before bony collapse of the lunate occurs***

# POST-TEST QUESTION #2

- ⦿ Definitive treatment for a Jersey finger injury...
  - A. is always conservative: 6-8 weeks of splinting typically does well.
  - B. may be conservative or surgical, it depends on the location of the injury.
  - C. is always surgical (primary tendon repair or fracture fragment repair). Long-term splinting is rarely an option.
  - D. is a corticosteroid injection at the site of injury.

# POST-TEST QUESTION #2

- ⦿ Definitive treatment for a Jersey finger injury...
  - A. is always conservative: 6-8 weeks of splinting typically does well.
  - B. may be conservative or surgical, it depends on the location of the injury.
  - C. ***is always surgical (primary tendon repair or fracture fragment repair). Long-term splinting is rarely an option.***
  - D. is a corticosteroid injection at the site of injury.

# POST-TEST QUESTION #3

- ◉ When evaluating a patient with a suspected skier's thumb injury...
  - A. it is best to obtain radiographs prior to assessing the UCL.
  - B. radiographs are not necessary - it is a clinical diagnosis.
  - C. it is best to obtain radiographs after assessing the UCL.
  - D. MRI is the gold standard imaging that is needed.

# POST-TEST QUESTION #3

- ◉ When evaluating a patient with a suspected skier's thumb injury...
  - A. *it is best to obtain radiographs prior to assessing the UCL.*
  - B. radiographs are not necessary - it is a clinical diagnosis.
  - C. it is best to obtain radiographs after assessing the UCL.
  - D. MRI is the gold standard imaging that is needed.

# CLINICAL CITATIONS

1. Blazer, PE. History and examination of the adult with hand pain. In: UpToDate, Curtis, MR (Ed), UpToDate, Waltham, MA, 2018
2. Boggess, BR. Evaluation of the adult with acute wrist pain. In: UpToDate, Grayzel, J (Ed), UpToDate, Waltham, MA, 2018
3. Boggess, BR. Evaluation of the adult with subacute or chronic wrist pain. In: UpToDate, Fields, KB (Ed), UpToDate, Waltham, MA, 2018
4. Bray, J and Neal, S. Evaluation of the patient with thumb pain. In: UpToDate, Grayzel, J (Ed), UpToDate, Waltham, MA, 2018
5. Miller, MD, Thompson, SR. Miller's review of orthopaedics. 6th ed. Philadelphia, PA: Elsevier; 2012.
6. Rynders, SD, Hart JA. Orthopedics for physician assistants. Philadelphia, PA: Elsevier; 2013.